OpenEvidence has signed content agreements with **NEJM**, **JAMA**, **NCCN**, **Wiley**, **Cochrane**, and more.

This conversation has been affirmatively and intentionally made public by the user who created it.

How likely is a DVT to develop in an austere setting, such as in prolonged field care or after a backcountry injury that leaves someone immobile while pending evacuation?

Finished Thinking

The risk of DVT in an austere setting with traumatic injury and prolonged immobilization is **substantial**, and the literature consistently places it among the highest-risk scenarios for VTE. The exact incidence depends heavily on injury pattern, duration of immobility, and whether any prophylaxis is available.

Incidence Estimates by Scenario

Without any prophylaxis, the landmark Geerts et al. (1994) prospective study found a **58% DVT rate** in severely injured trauma patients screened with venography. NEJM + 1 In patients with lower-limb fractures specifically, venographically-proven DVT rates range from **27% to 78%** depending on fracture location and severity. Cochrane Proximal fractures (femur, pelvis) carry the highest risk. A 2024 systematic review and meta-analysis of over 1.9 million trauma patients confirmed that lower extremity injury, pelvic injury, spinal injury, higher Injury Severity Score, older age, obesity, and male sex are all independently associated with increased VTE risk. Trauma and Ac...

For symptomatic VTE (the clinically relevant events), the American College of Chest Physicians estimates a baseline risk of **at least 3–5%** in most major trauma patients, rising to **8–10%** in those with traumatic brain or spinal cord injury. Chest In patients with lower-limb immobilization in a cast or brace for ≥1 week without prophylaxis, the Cochrane review found symptomatic DVT rates of **4.3% to 40%**. Cochrane

The following figure from the Geerts et al. study illustrates how DVT frequency varies by injury site and combination, with lower extremity injuries consistently driving the highest rates:



The NEW ENGLAND JOURNAL of MEDICINE



Figure 1. The Frequency of Deep-Vein Thrombosis among 349 Patients with One or More Major Injuries.

A Prospective Study of Venous Thromboembolism after Major Trauma.
N Engl J Med. December 15, 1994.
Used under license from The New England Journal of Medicine.

Timeline of Risk

DVT risk begins accruing rapidly after injury. One cohort study found that immobility for **more than 3 days** significantly increases the risk of proximal DVT compared to mobile patients (OR 3.59). Vascular Medici... The highest overall VTE incidence occurs in the **first 3 months** post-injury (10.3%), dropping to 0.5% by 6 months. Trauma and Ac... Critically, a delay

in pharmacologic prophylaxis of even **24–48 hours** independently confers a roughly **twofold increase** in VTE incidence. Trauma and Ac... + 1

Unique Risk Factors in Austere/Prolonged Field Care Settings

The Department of Defense Joint Trauma System Clinical Practice Guideline specifically identifies several factors that compound VTE risk in combat and austere environments: Military Medicine

- Hypercoagulable state induced by trauma itself
- Massive extremity injuries
- **Prolonged immobility** without access to mobilization
- **Long transport times** (fixed-wing evacuation, delayed evacuation)
- Dehydration and environmental exposure
- Tranexamic acid use (now identified as an independent VTE risk factor) Trauma and Ac...

These factors converge in a backcountry or prolonged field care scenario: the patient has sustained trauma, cannot ambulate, may be dehydrated, and faces hours to days before definitive care.

Prophylaxis Considerations in Resource-Limited Settings

In an austere environment where standard hospital-based prophylaxis (sequential compression devices, LMWH injections) may not be available, the following measures are relevant:

- **Mobilization** of uninjured extremities and early ambulation (even partial) when safe is the most accessible intervention, though its efficacy as sole prophylaxis is likely overstated. JAMA
- **Graduated compression stockings**, if available, may provide modest benefit, particularly when pharmacologic prophylaxis is not feasible. Trauma and Ac... + 1
- **LMWH** (e.g., enoxaparin 30 mg SQ q12h or 40 mg SQ daily) remains the standard pharmacologic agent and is shelf-stable, making it a reasonable addition to prolonged field care kits. Unfractionated heparin (5,000 U SQ q8–12h) is an alternative. Trauma and Ac... + 2
- **Aspirin** has been considered as a more logistically feasible option for extended prophylaxis post-discharge, though evidence in the acute trauma setting is limited. Trauma and Ac... + 1
- **Hydration** is reasonable to maintain, though direct evidence linking dehydration to VTE prevention is lacking. CDC Yellow Book

In summary, an immobilized trauma patient in a backcountry or prolonged field care setting faces a clinically significant DVT risk that may approach **50–60% for asymptomatic DVT** with major lower extremity trauma and no prophylaxis, or **3–10% for symptomatic VTE** depending on injury severity. This risk begins within the first days and escalates with each day of immobility and delayed prophylaxis.

Would you like to explore specific prolonged field care protocols for VTE prophylaxis, including practical considerations for carrying and administering LMWH in austere environments?

References

1. A Prospective Study of Venous Thromboembolism after Major Trauma.
   The New England Journal of Medicine. 1994. Geerts WH, Code KI, Jay RM, Chen E, Szalai JP.

2. Updated Guidelines to Reduce Venous Thromboembolism in Trauma Patients: A Western Trauma Association Critical Decisions Algorithm.
   The Journal of Trauma and Acute Care Surgery. 2020. Ley EJ, Brown CVR, Moore EE, et al. Guideline

3. Low Molecular Weight Heparin for Prevention of Venous Thromboembolism in Patients With Lower-Limb Immobilization.
   The Cochrane Database of Systematic Reviews. 2017. Zee AA, van Lieshout K, van der Heide M, Janssen L, Janzing HM.

4. Prognostic Factors Associated With Venous Thromboembolism Following Traumatic Injury: A Systematic Review and Meta-Analysis.
   The Journal of Trauma and Acute Care Surgery. 2024. Tran A, Fernando SM, Rochwerg B, et al.

5. Prevention of VTE in Nonorthopedic Surgical Patients: Antithrombotic Therapy and Prevention of Thrombosis, 9th Ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines.
   Chest. 2012. Gould MK, Garcia DA, Wren SM, et al. Guideline

6. Relevance of Immobility as a Risk Factor for Symptomatic Proximal and Isolated Distal Deep Vein Thrombosis in Acutely Ill Medical Inpatients.
   Vascular Medicine. 2021. Sartori M, Favaretto E, Cosmi B.

7. Can We Ever Stop Worrying About Venous Thromboembolism After Trauma?.
   The Journal of Trauma and Acute Care Surgery. 2015. Godat LN, Kobayashi L, Chang DC, Coimbra R.

8. Timing of Venous Thromboembolism Prophylaxis Initiation After Injury: Findings From the Consensus Conference to Implement Optimal VTE Prophylaxis in Trauma.
   The Journal of Trauma and Acute Care Surgery. 2023. Schellenberg M, Costantini T, Joseph B, et al.

9. Prevention of Deep Venous Thromboembolism.
Military Medicine. 2018. Grabo DJ, Seery JM, Bradley M, et al.

10. Venous Thromboembolism Prevention in Emergency General Surgery: A Review.
JAMA Surgery. 2018. Murphy PB, Vogt KN, Lau BD, et al.

11. Deep Vein Thrombosis and Pulmonary Embolism.
CDC Yellow Book. 2025. Nimia L. Reyes and Karon Abe

12. Prophylaxis of Venous Thromboembolism in Trauma: What You Need to Know.
The Journal of Trauma and Acute Care Surgery. 2026. Lippincott M, Berndtson AE. New

13. Approaches for Optimizing Venous Thromboembolism Prevention in Injured Patients: Findings From the Consensus Conference to Implement Optimal Venous Thromboembolism Prophylaxis in Trauma.
The Journal of Trauma and Acute Care Surgery. 2023. Teichman AL, Cotton BA, Byrne J, et al.

Follow-Up Questions

What are the most effective DVT prophylaxis options in resource-limited environments? ›

Which patient populations in austere settings are at highest risk for DVT development? ›

What are the potential complications of untreated DVT in such scenarios? ›